

**UNCLASSIFIED**

---

**AD 400 348**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-3-1

IMM-NYU 307  
JANUARY 1963

CATALOGED BY ASTIA  
AS AD NO. 480 348

400 348



NEW YORK UNIVERSITY  
COURANT INSTITUTE OF  
MATHEMATICAL SCIENCES

# The Solution of Some Non-Linear Integral Equations with Cauchy Kernels

A. S. PETERS

ASTIA  
APR 5 1963

---

PREPARED UNDER  
CONTRACT NO. NONR-285(06)  
WITH THE  
OFFICE OF NAVAL RESEARCH

New York University  
Courant Institute of Mathematical Sciences

THE SOLUTION OF SOME NON-LINEAR INTEGRAL EQUATIONS  
WITH CAUCHY KERNELS

A. S. Peters

This report represents results obtained at the  
Courant Institute of Mathematical Sciences,  
New York University, with the Office of Naval  
Research, Contract No. Nonr-285(06).  
Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

## 1. Introduction

In the theory of radiative transfer there are several problems which can be solved by finding the solution  $H(u)$  of the non-linear integral equation

$$(1.1) \quad \frac{1}{H(x)} = 1 - x \int_0^1 \frac{g(u)H(u)du}{u+x}$$

which is called Chandrasekhar's equation. The books by Chandrasekhar [1] and Kourganoff [2] contain discussions of this important equation; and these books also present the contributions of various mathematicians who have shown that (1.1) can be solved explicitly by using function theory techniques based on analytic continuation. Recently, (1961) C. Fox [3] has shown that (1.1) can be converted into the linear equation

$$(1.2) \quad H(x)G(x) = 1 + x \int_0^1 \frac{g(u)H(u)du}{u-x}$$

where  $G(x)$  is known and  $g(x)$  is prescribed. The equation (1.2) is a singular integral equation with a Cauchy kernel and it can be solved for  $H(u)$  by using an extension of Carleman's method as shown for example in Muskhelishvili's book [4].

Chandrasekhar's equation can be linearized by first writing it in the form

$$(1.3) \quad xg(x) = \lambda\phi_1(x) + \phi_1(x) \int_0^1 \frac{\phi_1(u)du}{u+x}$$

and then integrating each side of this equation after it has been multiplied by the factor  $1/(x+\xi)$ . This gives

$$(1.4) \quad \int_0^1 \frac{xg(x)du}{x+\xi} = \left[ \lambda + \int_0^1 \frac{\phi_1(u)du}{u-\xi} \right] \int_0^1 \frac{\phi_1(x)dx}{x+\xi} \\ - \int_0^1 \frac{\phi_1(u)}{u-\xi} \int_0^1 \frac{\phi_1(x)dx}{x+u} du$$

and after multiplying (1.4) by  $\phi_1(\xi)$  and using (1.3) we find

$$(1.5) \quad \phi_1(\xi)G_1(\xi) = \lambda + \int_0^1 \frac{\phi_1(x)dx}{x-\xi}.$$

This is essentially the procedure that was used by Fox to pass from (1.1) to (1.2). It suggests the possibility of solving

$$(1.6) \quad \lambda\phi_1(x) + \phi_1(x) \int_0^1 \frac{\phi_1(u)du}{u-x} = f_1(x)$$

which is both singular and non-linear. Equation (1.6), in turn, suggests an investigation of the more general equation

$$I \quad \lambda\phi(\zeta) + \phi(\zeta) \int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} = f(\zeta)$$

where  $\zeta$  is an interior point of the simple smooth arc  $L$  which connects the points  $\tau_0$  and  $\tau_1$  in the complex  $\tau$ -plane.

One of the purposes of this paper is to show in Section 2 that the equation I can be solved explicitly by using elementary function theory techniques. It turns out that the solution of I is in some ways simpler than the solution of (1.3). We will also be concerned with the solution of some other non-linear equations. In Section 3 we show how to solve

$$\text{II} \quad \pi^2 \phi^2(\zeta) - \left[ \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} \right]^2 = f(\zeta)$$

and Section 4 is devoted to the solution of

$$\text{III} \quad \pi^2 \phi^2(\zeta) + \left[ \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} \right]^2 = f(\zeta) .$$

In Section 5 we show that there is a connection between equations I, II, III and certain problems in potential theory.

The equations I, II, III may be of interest for at least two reasons. In the first place, they are of interest in themselves as Cauchy singular, non-linear integral equations which can be solved explicitly. In the second place, they present a formulation of certain non-linear boundary value problems. Equation III, for example, is intimately related to a problem in two-dimensional potential theory which has a number of physical applications. This is the problem of finding a potential function in a domain  $D$  when its normal derivative is prescribed on one part of the boundary  $C$ ; and the magnitude of its gradient is given on the remaining part of  $C$ . In Section 5 we show how an explicit formula for the solution of this problem can be found.

The final Section 6 is concerned with a brief discussion of the non-linear system

$$\phi(\zeta) \int_L \frac{\psi(\tau) d\tau}{\tau - \zeta} = f(\zeta)$$

$$\psi(\zeta) \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} = g(\zeta)$$

and some other systems which can be linearized by the method developed in Section 2.

We state here the main conditions and assumptions upon which our analysis is based. If  $\tau = \tau(t)$  is the equations of the simple smooth arc  $L$  directed from  $\tau_0$  to  $\tau_1 \neq \tau_0$  let  $L[\tau_0, \tau_1]$  denote the set of points  $\tau = \tau(t)$ ,  $t_0 \leq t \leq t_1$ ; and let  $L(\tau_0, \tau_1)$  denote the set  $L[\tau_0, \tau_1]$  minus the endpoints  $\tau_0$  and  $\tau_1$ . We assume for simplicity that the unknown function  $\phi(\tau)$  satisfies a uniform Hölder condition for any pair of points in  $L(\tau_0, \tau_1)$  and thus guarantee the existence of the Cauchy principal values of the integrals which appear in I, II, III where  $\zeta$  is in  $L(\tau_0, \tau_1)$ . We also assume that if  $\phi(\tau)$  has a singularity at an endpoint  $\alpha$  of  $L[\tau_0, \tau_1]$  it is such that  $\lim_{\tau \rightarrow \alpha} (\tau - \alpha)\phi(\tau) = 0$ . We will say that a function with these properties belongs to the class  $\mathcal{H}_1$ . If  $\phi(\tau)$  belongs to the class  $\mathcal{H}_1$  it can be shown that

$$\int_L \frac{\phi(\tau) d\tau}{\tau - \zeta}$$

satisfies a uniform Hölder condition for any pair of points  $\zeta_1$  and  $\zeta_2$  in  $L(\tau_0, \tau_1)$ ; and

$$F(z) = \int_L \frac{\phi(\tau) d\tau}{\tau - z}$$

as a function of the complex variable  $z$  is such that

$$\lim_{z \rightarrow \alpha} (z - \alpha)F(z) = 0.$$

The properties of the prescribed function must match the properties implied by the representations on the left-hand side of I, II, III. That is, in accordance with our assumptions about



$\phi(\tau)$ ,  $f(\zeta)$  must satisfy a uniform Hölder condition on  $L(\tau_0, \tau_1)$ ; and although it may have a stronger singularity than  $\phi(\tau)$  at  $\alpha$ ,  $f(\zeta)$  must be such that  $\lim_{\zeta \rightarrow \alpha} (\zeta - \alpha)^2 f(\zeta) = 0$ . We will say that under these conditions  $f(\zeta)$  belongs to the class  $\mathcal{A}_2$ .

The transformation  $2\tau = (\tau_1 - \tau_0)v + (\tau_0 + \tau_1)$  maps  $L[\tau_0, \tau_1]$  into  $L[-1, 1]$  a simple smooth arc directed from  $\tau = -1$  to  $\tau = 1$ . If we also use  $2\zeta = (\tau_1 - \tau_0)\omega + (\tau_0 + \tau_1)$  we have

$$\int_{L[\tau_0, \tau_1]} \frac{\phi(\tau) d\tau}{\tau - \zeta} = \int_{L[-1, 1]} \frac{\phi_0(v) dv}{v - \omega}$$

which shows that the transformation does not change the form of equations I, II, III. Thus there is no loss of generality if we assume, as we will hereafter, that  $L$  in I, II, III is  $L[-1, 1]$ .

## 2. Equation I

We proceed to show how

$$(2.1) \quad \lambda \phi(\zeta) + \phi(\zeta) \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} = f(\zeta)$$

can be solved by using the Hardy-Poincaré-Bertrand formula. This formula states that if  $\psi(\tau, \zeta)$  is suitably restricted then

$$(2.2) \quad \int_L \frac{1}{\tau - \omega} \int_L \frac{\psi(\tau, \zeta) d\tau d\zeta}{\tau - \zeta} = -\pi^2 \psi(\omega, \omega) + \int_L \int_L \frac{\psi(\tau, \zeta) d\zeta d\tau}{(\tau - \zeta)(\zeta - \omega)}$$

provided  $\omega$  is in  $L(-1, 1)$ . This says that an interchange of the order of integration on the left side of (2.2) leaves the residue  $-\pi^2 \psi(\omega, \omega)$ . If  $\phi(\tau)$  belongs to the class  $\mathcal{A}_1$ , defined in the

introduction, the formula (2.2) holds with  $\psi(\tau, \zeta) = (1-\zeta^2)\phi(\zeta)\phi(\tau)$ .

The application of (2.2) to

$$\frac{1}{1-\omega^2} \int_L \frac{(1-\zeta^2)\phi(\zeta)}{\zeta-\omega} \int_L \frac{\phi(\tau)d\tau d\zeta}{\tau-\zeta}$$

shows, after a little manipulation, that

$$(2.3) \quad \frac{2}{1-\omega^2} \int_L \frac{(1-\zeta^2)\phi(\zeta)}{\zeta-\omega} \int_L \frac{\phi(\tau)d\tau d\zeta}{\tau-\zeta} \\ = -\pi^2 \phi^2(\omega) + \left[ \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} \right]^2 + \frac{1}{1-\omega^2} \left[ \int_L \phi(\tau)d\tau \right]^2.$$

Now if a solution of (2.1) exists when  $f(\zeta)$  belongs to  $\mathcal{A}_2$ , then (2.1) and (2.3) imply

$$(2.4) \quad \frac{2}{1-\omega^2} \int_L \frac{(1-\zeta^2)f(\zeta)d\zeta}{\zeta-\omega} = \frac{2\lambda}{1-\omega^2} \int_L \frac{(1-\zeta^2)\phi(\zeta)d\zeta}{\zeta-\omega} - \pi^2 \phi^2(\omega) \\ + \left[ \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} \right]^2 + \frac{1}{1-\omega^2} \left[ \int_L \phi(\tau)d\tau \right]^2$$

or

$$(2.5) \quad \frac{2}{1-\omega^2} \int_L \frac{(1-\zeta^2)f(\zeta)d\zeta}{\zeta-\omega} = -\pi^2 \phi^2(\omega) + 2\lambda \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} \\ + \left[ \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} \right]^2 + \frac{[k_0^2 - 2\lambda(k_1 + \omega k_0)]}{1-\omega^2}$$

where

$$k_0 = \int_L \phi(\tau)d\tau$$

$$k_1 = \int_L \tau \phi(\tau)d\tau.$$

The addition of

$$(2.6) \quad \pm 2\pi i f(\omega) = \pm \left[ 2\lambda\pi i \phi(\omega) + 2\pi i \phi(\omega) \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} \right]$$

to (2.5) gives

$$(2.7) \quad \frac{2}{1-\omega^2} \int_L \frac{(1-\zeta^2)f(\zeta)d\zeta}{\zeta-\omega} \pm 2\pi i f(\omega) \\ = \left[ \pm \pi i \phi(\omega) + \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} + \lambda \right]^2 - \lambda^2 + \frac{[k_0^2 - 2\lambda(k_1 + \omega k_0)]}{1-\omega^2}.$$

Let us introduce the function

$$F_0(z) = (1-z^2) \left[ \int_L \frac{\phi(\tau)d\tau}{\tau-z} + \lambda \right]^2 \\ - \left[ 2 \int_L \frac{(1-\tau^2)f(\tau)d\tau}{\tau-z} + 2\lambda(k_1 + zk_0) - k_0^2 + (1-z^2)\lambda^2 \right].$$

This function is analytic for  $z$  not on  $L[-1,1]$  and it vanishes as  $z \rightarrow \infty$ . As  $z$  approaches the point  $\omega$  from the positive and negative sides of  $L$ , the limit values of  $F_0(z)$  are

$$F_0^\pm(\omega) = (1-\omega^2) \left[ \pm \pi i \phi(\omega) + \int_L \frac{\phi(\tau)d\tau}{\tau-\omega} + \lambda \right]^2 \\ - \left[ \pm 2\pi i (1-\omega^2)f(\omega) + 2 \int_L \frac{(1-\tau^2)f(\tau)d\tau}{\tau-\omega} \right] \\ - [2\lambda(k_1 + \omega k_0) - k_0^2 + (1-\omega^2)\lambda^2]$$

but, as we can see from (2.7), these limit values vanish. Furthermore, it follows from the conditions we have imposed on  $\phi$  and  $f$ , that the behavior of  $F_0(z)$  in the neighborhood of an endpoint of  $L$  must be such that

$$(2.8) \quad \begin{aligned} \lim_{z \rightarrow 1} (1-z)F_0(z) &= 0 \\ \lim_{z \rightarrow -1} (1+z)F_0(z) &= 0. \end{aligned}$$

Let  $\Gamma$  be the boundary of a domain which contains  $L[-1,1]$ . If  $z$  is in the exterior of  $\Gamma$  then

$$F_0(z) = \frac{1}{2\pi i} \oint_{\Gamma} \frac{F_0(\xi)d\xi}{\xi - z}.$$

Since (2.8) holds, the path  $\Gamma$  can be collapsed into the path  $C$ , composed of the upper and lower banks of  $L$ , without changing the value of the Cauchy integral. At these banks, however, the limit values  $F_0^\pm(\omega)$  vanish. We therefore conclude that  $F_0(z)$  vanishes everywhere in the exterior of  $L[-1,1]$  and hence, if a solution of (2.1) exists, the constants  $k_0$  and  $k_1$  must be such that the two-valued function

$$(2.9) \quad S(z) = \left[ \frac{1}{1-z^2} \left\{ 2 \int_L \frac{(1-\tau^2)f(\tau)d\tau}{\tau-z} + 2\lambda(k_1 + zk_0) - k_0^2 + (1-z^2)\lambda^2 \right\} \right]^{\frac{1}{2}}$$

$$(A) \quad = \int_L \frac{\phi(\tau)d\tau}{\tau-z} + \lambda$$

is analytic for  $z$  not on  $L[-1,1]$ . We will refer to this condition as condition (A).

If for a prescribed  $f(\zeta)$  constants  $k_0$  and  $k_1$  exist such that  $S(z)$  is analytic for  $z$  not on  $L[-1,1]$ , then, as we can see by taking the limit values of (2.9),

$$(2.10) \quad \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} + \pi i \phi(\zeta) + \lambda = S^+(\zeta)$$

$$(2.11) \quad \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} - \pi i \phi(\zeta) + \lambda = S^-(\zeta)$$

and, as subtraction of (2.11) from (2.10) shows, the solution of (2.1) is

$$\begin{aligned} \phi(\zeta) &= \frac{1}{2\pi i} [S^+(\zeta) - S^-(\zeta)] \\ (2.12) \quad &= \frac{1}{2\pi i} \left[ \frac{2}{1-\zeta^2} \int_L \frac{(1-\tau^2)f(\tau)d\tau}{\tau-\zeta} + 2\pi i f(\zeta) + \lambda^2 \right]^{\frac{1}{2}} \\ &\quad + \frac{2\lambda(k_1 + \zeta k_0) - k_0^2}{1-\zeta^2} \\ &\quad - \frac{1}{2\pi i} \left[ \frac{2}{1-\zeta^2} \int_L \frac{(1-\tau^2)f(\tau)d\tau}{\tau-\zeta} - 2\pi i f(\zeta) + \lambda^2 \right]^{\frac{1}{2}} \\ &\quad + \frac{2\lambda(k_1 + \zeta k_0) - k_0^2}{1-\zeta^2} \end{aligned}$$

For the case in which  $L[-1,1]$  coincides with the real axis we find that if  $f(\xi)$  is real then the solution of

$$(2.13) \quad \lambda \phi(\xi) + \phi(\xi) \int_{-1}^1 \frac{\phi(t) dt}{t - \xi} = f(\xi), \quad -1 < \xi < 1$$

is

$$(2.14) \quad \phi(\xi) = \frac{1}{\pi} \mathcal{Q} \left[ \frac{2}{1-\xi^2} \int_{-1}^1 \frac{(1-t^2)f(t)dt}{t-\xi} + 2\pi i f(\xi) + \lambda^2 \right]^{\frac{1}{2}} \\ + \frac{1}{1-\xi^2} \{2\lambda(k_1 + \xi k_0) - k_0^2\}$$

It is easy to verify that (2.12) satisfies (2.1) when  $S(z)$  satisfies condition (A). If we write

$$\phi(\zeta) = \frac{1}{2\pi i} [S_0(\zeta) + 2\pi i f(\zeta)]^{\frac{1}{2}} \\ - \frac{1}{2\pi i} [S_0(\zeta) - 2\pi i f(\zeta)]^{\frac{1}{2}}$$

we have

$$\int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} = \frac{1}{2\pi i} \int_L \frac{\{[S_0(\tau) + 2\pi i f(\tau)]^{\frac{1}{2}} - [S_0(\tau) - 2\pi i f(\tau)]^{\frac{1}{2}}\}d\tau}{\tau-\zeta}, \\ \int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} = \frac{1}{2\pi i} \oint_C \frac{S(\tau)d\tau}{\tau-\zeta}.$$

Now if we detach  $C$  from  $L$  and expand  $C$  into  $\Gamma$  we have

$$\int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} = \frac{1}{2} \left\{ [S_0(\zeta) + 2\pi i f(\zeta)]^{\frac{1}{2}} + [S_0(\zeta) - 2\pi i f(\zeta)]^{\frac{1}{2}} \right\} \\ + \frac{1}{2\pi i} \oint_{\Gamma} \frac{S(\tau)d\tau}{\tau-\zeta}$$

and, since  $S(\tau) \rightarrow \lambda$  as  $\tau \rightarrow \infty$ ,

$$\int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} = \frac{1}{2} \left\{ [S_0(\zeta) + 2\pi i f(\zeta)]^{\frac{1}{2}} + [S_0(\zeta) - 2\pi i f(\zeta)]^{\frac{1}{2}} \right\} - \lambda.$$

Therefore

$$\begin{aligned}
\lambda \phi(\zeta) + \phi(\zeta) \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} &= \frac{\lambda}{2\pi i} [S^+(\zeta) - S^-(\zeta)] \\
&+ \frac{1}{4\pi i} [S_0(\zeta) + 2\pi i f(\zeta) - S_0(\zeta) + 2\pi i f(\zeta)] \\
&- \frac{\lambda}{2\pi i} [S^+(\zeta) - S^-(\zeta)] \\
&= f(\zeta) .
\end{aligned}$$

The condition that  $S(z)$  must be analytic in the complex plane slit along  $L[-1,1]$ , [the condition (A)], does not in general determine the constants  $k_0$  and  $k_1$  numerically. This can be seen by considering the equation

$$(2.15) \quad \lambda \phi(\xi) + \phi(\xi) \int_{-1}^1 \frac{\phi(t) dt}{t - \xi} = 0 .$$

The function  $S(z)$  for this equation is

$$\begin{aligned}
S(z) &= \left[ \lambda^2 + \frac{k_0^2 - 2\lambda k_1 - 2\lambda k_0 z}{z^2 - 1} \right]^{\frac{1}{2}} \\
&= \left[ \frac{\lambda^2 z^2 - 2\lambda k_0 z + k_0^2 - 2\lambda k_1 - \lambda^2}{z^2 - 1} \right]^{\frac{1}{2}}
\end{aligned}$$

and this certainly satisfies condition (A) if we take  $k_0 = k_1 = 0$ . For these values we have

$$\phi(\xi) = \frac{1}{\pi} \int (\lambda^2)^{\frac{1}{2}} = 0 ,$$

namely the trivial solution of (2.15). On the other hand,  $S(z)$  will satisfy (A) if

$$\lambda^2 z^2 - 2\lambda k_0 z + k_0^2 - 2\lambda k_1 - \lambda^2$$

possesses a double zero, that is, if we take  $2k_1 + \lambda = 0$ . With this choice,  $k_0$  is arbitrary, and

$$S(z) = \frac{\lambda z - k_0}{\sqrt{z^2 - 1}}$$

from which

$$\phi(\xi) = \frac{1}{\pi} \oint \frac{\lambda \xi - k_0}{\sqrt{\xi^2 - 1}} = \frac{1}{\pi} \cdot \frac{\lambda \xi - k_0}{\sqrt{1 - \xi^2}}$$

and this is the general solution of

$$\lambda + \int_{-1}^1 \frac{\phi(t) dt}{t - \xi} = 0.$$

Thus we can see that in order to fix  $k_0$  and  $k_1$  uniquely we may have to impose side conditions in addition to condition (A).

Let us examine the possibility of having a solution  $\phi$  whose endpoint behavior is such that

$$(B) \quad \mathcal{L}_{\tau \rightarrow \alpha} (\alpha - \tau)^{\frac{1}{2}} \phi(\tau) = 0.$$

With this behavior the endpoint behavior of  $f(\zeta)$  must be such that

$$(B_1) \quad \mathcal{L}_{\zeta \rightarrow \alpha} (\alpha - \zeta) f(\zeta) = 0.$$

Under condition (B) the order of integration can be changed in

$$\int_L \phi(\zeta) \int_L \frac{\phi(\tau) d\tau d\zeta}{\tau - \zeta}$$

and hence by integrating (2.1) we find



$$\lambda \int_L \phi(\zeta) d\zeta - \int_L \phi(\tau) \int_L \frac{\phi(\zeta) d\zeta d\tau}{\zeta - \tau} = \int_L f(\zeta) d\zeta$$

$$\lambda \int_L \phi(\zeta) d\zeta + \int_L [\lambda \phi(\tau) - f(\tau)] d\tau = \int_L f(\zeta) d\zeta$$

$$(2.16) \quad \lambda k_0 = \int_L f(\tau) d\tau .$$

Also, after multiplying (2.1) by  $\zeta$  and integrating:

$$\lambda \int_L \zeta \phi(\zeta) d\zeta - \int_L \phi(\tau) \int_L \frac{(\zeta - \tau + \tau) \phi(\zeta) d\zeta d\tau}{\zeta - \tau} = \int_L \zeta f(\zeta) d\zeta$$

$$\lambda \int_L \zeta \phi(\zeta) d\zeta - \left[ \int_L \phi(\tau) d\tau \right]^2 + \int_L \tau [\lambda \phi(\tau) - f(\tau)] d\tau = \int_L \zeta f(\zeta) d\zeta$$

$$(2.17) \quad 2\lambda k_1 - 2 \int_L \tau f(\tau) d\tau = k_0^2 .$$

The substitution of these values for  $k_0$  and  $k_1$  in (2.9) gives

$$(2.18) \quad S(z) = \left[ 2 \int_L \frac{f(\tau) d\tau}{\tau - z} + \lambda^2 \right]^{\frac{1}{2}}$$

and a solution satisfying (B) will exist if  $f(\zeta)$  satisfies (B<sub>1</sub>) and (2.18) satisfies (A). When this is the case the solution is

$$\begin{aligned} \phi(\zeta) = & \frac{1}{2\pi i} \left[ 2 \int_L \frac{f(\tau) d\tau}{\tau - \zeta} + 2\pi i f(\zeta) + \lambda^2 \right]^{\frac{1}{2}} \\ & - \frac{1}{2\pi i} \left[ 2 \int_L \frac{f(\tau) d\tau}{\tau - \zeta} - 2\pi i f(\zeta) + \lambda^2 \right]^{\frac{1}{2}} . \end{aligned}$$

Instead of imposing the condition (B) at each endpoint of  $L$  we may wish to impose it at only one endpoint, say the condition

$$(C) \quad \mathcal{L}_{\tau \rightarrow 1} (1-\tau)^{\frac{1}{2}} \phi(\tau) = 0.$$

The function  $f(\zeta)$  would then have to be such that

$$(C_1) \quad \mathcal{L}_{\zeta \rightarrow 1} (1-\zeta)f(\zeta) = 0.$$

Under condition (C) we find from (2.1) that

$$\begin{aligned} \lambda \int_L (1+\zeta)\phi(\zeta)d\zeta - \int_L \phi(\tau) \int_L \frac{(\zeta-\tau+1)\phi(\zeta)d\zeta d\tau}{\zeta-\tau} &= \int_L (1+\zeta)f(\zeta)d\zeta \\ \lambda \int_L (1+\zeta)\phi(\zeta)d\zeta - \left[ \int_L \phi(\tau)d\tau \right]^{\frac{1}{2}} + \int_L (1+\tau)[\lambda\phi(\tau)-f(\tau)]d\tau &= \int_L (1+\zeta)f(\zeta)d\zeta \\ (2.19) \quad 2\lambda(k_0 + k_1) - k_0^2 &= 2 \int_L (1+\tau)f(\tau)d\tau. \end{aligned}$$

The substitution of this value of  $k_1$  in (2.9) gives

$$(2.20) \quad S(z) = \left[ \frac{2}{1+z} \int_L \frac{(1+\tau)f(\tau)d\tau}{\tau-z} - \frac{2\lambda k_0}{1+z} + \lambda^2 \right]^{\frac{1}{2}}$$

and a solution satisfying (C) will exist if  $f(\zeta)$  satisfies  $(C_1)$  and (2.20) satisfies (A). When this is the case the solution is

$$\begin{aligned} \phi(\zeta) &= \frac{1}{2\pi i} \left[ \frac{2}{1+\zeta} \int_L \frac{(1+\tau)f(\tau)d\tau}{\tau-\zeta} + 2\pi i f(\zeta) - \frac{2\lambda k_0}{1+\zeta} + \lambda^2 \right]^{\frac{1}{2}} \\ &\quad - \frac{1}{2\pi i} \left[ \frac{2}{1+\zeta} \int_L \frac{(1+\tau)f(\tau)d\tau}{\tau-\zeta} - 2\pi i f(\zeta) - \frac{2\lambda k_0}{1+\zeta} + \lambda^2 \right]^{\frac{1}{2}}. \end{aligned}$$

### 3. Equation II

In this section, we show that

$$(3.1) \quad \pi^2 \phi^2(\zeta) - \left[ \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} \right]^2 = f(\zeta)$$

can be converted into a particular case of (2.1). We saw in Section 2 that if  $\phi(\tau)$  belongs to the class  $\mathcal{A}_1$  then we have the identity

$$(3.2) \quad \frac{2}{1-\omega^2} \int_L \frac{(1-\zeta^2)\phi(\zeta)}{\zeta-\omega} \int_L \frac{\phi(\tau) d\tau d\zeta}{\tau-\zeta} \\ = -\pi^2 \phi^2(\omega) + \left[ \int_L \frac{\phi(\tau) d\tau}{\tau-\omega} \right]^2 + \frac{1}{1-\omega^2} \left[ \int_L \phi(\tau) d\tau \right]^2.$$

Hence if  $\phi(\tau)$  satisfies (3.1) then it must satisfy

$$(3.3) \quad \frac{2}{1-\omega^2} \int_L \frac{(1-\zeta^2)\phi(\zeta)}{\zeta-\omega} \int_L \frac{\phi(\tau) d\tau d\zeta}{\tau-\zeta} = -f(\omega) + \frac{k_0^2}{1-\omega^2}$$

or

$$(3.4) \quad 2 \int_L \frac{(1-\zeta^2)\phi(\zeta)}{\zeta-\omega} \int_L \frac{\phi(\tau) d\tau d\zeta}{\tau-\zeta} = -(1-\omega^2)f(\omega) + k_0^2$$

where

$$k_0 = \int_L \phi(\tau) d\tau.$$

Equation (3.4) is a singular integral equation of the first kind which can be solved for

$$2(1-\zeta^2)\phi(\zeta) \int_L \frac{\phi(\tau) d\tau}{\tau-\zeta}.$$

The solution is

$$(3.5) \quad 2(1-\zeta^2)\phi(\zeta) \int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} \\ = - \frac{1}{\pi^2 \sqrt{1-\zeta^2}} \int_L \frac{\sqrt{1-\tau^2} [k_0^2 - (1-\tau^2)f(\tau)]d\tau}{\tau-\zeta} + \frac{k}{\sqrt{1-\zeta^2}}$$

where

$$\pi k = 2 \int_L (1-\zeta^2)\phi(\zeta) \int_L \frac{\phi(\tau)d\tau d\zeta}{\tau-\zeta} \\ = -2 \int_L (1-\tau^2)\phi(\tau) \int_L \frac{\phi(\zeta)d\zeta d\tau}{\zeta-\tau} + 2 \int_L \phi(\tau) \int_L (\zeta+\tau)\phi(\zeta)d\zeta d\tau$$

and therefore

$$\pi k = \int_L \phi(\tau) \int_L (\zeta+\tau)\phi(\zeta)d\zeta d\tau$$

$$(3.6) \quad \pi k = 2k_0 k_1 .$$

From (3.5) and (3.6) we find that if  $\phi(\tau)$  satisfies (3.1) it must satisfy

$$(3.7) \quad \phi(\zeta) \int_L \frac{\phi(\tau)d\tau}{\tau-\zeta} = F(\zeta)$$

where

$$(3.8) \quad F(\zeta) = \frac{1}{2\pi^2(1-\zeta^2)^{3/2}} \int_L \frac{(1-\tau^2)^{3/2}f(\tau)d\tau}{\tau-\zeta} \\ + \frac{\zeta k_0^2}{2\pi(1-\zeta^2)^{3/2}} + \frac{k_0 k_1}{\pi(1-\zeta^2)^{3/2}} .$$

Conversely, if  $\phi(\tau)$  satisfies (3.7) it must satisfy (3.1).

Equation (3.7) is equation (2.1) with  $\lambda = 0$ . We can apply the theory of Section 2. It follows from the results of Section 2 that (3.1) possesses a solution provided

$$S_1(z) = \left[ \frac{1}{(1-z^2)} \int_L \frac{(1-\tau^2)F(\tau)d\tau}{\tau-z} - \frac{k_0^2}{1-z^2} \right]^{\frac{1}{2}}$$

satisfies condition (A), i.e.,  $S_1(z)$  is analytic for  $z$  not on  $L$ .

Since

$$\begin{aligned} \frac{2}{1-z^2} \int_L \frac{(1-\tau^2)F(\tau)d\tau}{\tau-z} &= \frac{1}{\pi^2(1-z^2)} \int_L \frac{1}{(\tau-z)\sqrt{1-\tau^2}} \int_L \frac{(1-\zeta^2)^{3/2}f(\zeta)d\zeta}{\zeta-\tau} \\ &\quad + \frac{k_0^2}{\pi(1-z^2)} \int_L \frac{\tau d\tau}{(\tau-z)\sqrt{1-\tau^2}} + \frac{2k_0k_1}{\pi(1-z^2)} \int_L \frac{d\tau}{(\tau-z)\sqrt{1-\tau^2}} \\ &= \frac{1}{\pi(1-z^2)^{3/2}} \int_L \frac{(1-\zeta^2)^{3/2}f(\zeta)d\zeta}{\zeta-z} \\ &\quad + \frac{k_0^2}{(1-z^2)} + \frac{1k_0(2k_1+k_0z)}{(1-z^2)^{3/2}} \end{aligned}$$

we see that if (3.1) possesses a solution then the constants  $k_0$  and  $k_1$  must be such that

$$S_1(z) = \left[ \frac{1}{\pi(1-z^2)^{3/2}} \int_L \frac{(1-\zeta^2)^{3/2}f(\zeta)d\zeta}{\zeta-z} + \frac{1k_0(2k_1+k_0z)}{(1-z^2)^{3/2}} \right]^{\frac{1}{2}}$$

is analytic for  $z$  in the exterior of  $L$ . If this condition can be satisfied by a proper choice of  $k_0$  and  $k_1$ , the solution of (3.1) is

$$\begin{aligned}
 (3.9) \quad \phi(\tau) = & \frac{1}{2\pi i} \left[ -f(\tau) + \frac{1}{\pi^2(1-\tau^2)^{3/2}} \int_L \frac{(1-\xi^2)^{3/2} f(\xi) d\xi}{\xi - \tau} \right. \\
 & \left. + \frac{1k_0(2k_1 + k_0\tau)}{(1-\tau^2)^{3/2}} \right]^{\frac{1}{2}} \\
 & - \frac{1}{2\pi i} \left[ -f(\tau) - \frac{1}{\pi^2(1-\tau^2)^{3/2}} \int_L \frac{(1-\xi^2)^{3/2} f(\xi) d\xi}{\xi - \tau} \right. \\
 & \left. - \frac{1k_0(2k_1 + k_0\tau)}{(1-\tau^2)^{3/2}} \right]^{\frac{1}{2}}
 \end{aligned}$$

For the case in which  $L[-1,1]$  coincides with the real axis we find that if  $f(\xi)$  is real then the real solution of

$$(3.10) \quad \pi^2 \phi^2(\xi) - \left[ \int_{-1}^1 \frac{\phi(t) dt}{t - \xi} \right]^2 = f(\xi), \quad -1 < \xi < 1$$

is

$$(3.11) \quad \phi(\xi) = \frac{1}{\pi} \left[ -f(\xi) + \frac{1}{\pi(1-\xi^2)^{3/2}} \int_{-1}^1 \frac{(1-t^2)^{3/2} f(t) dt}{t - \xi} \right. \\
 \left. + \frac{1k_0(2k_1 + k_0\xi)}{(1-\xi^2)^{3/2}} \right]^{\frac{1}{2}}$$

provided condition (A) is satisfied.

#### 4. Equation III

The equation

$$(4.1) \quad \pi^2 \phi^2(\xi) + \left[ \int_L \frac{\phi(\tau) d\tau}{\tau - \xi} \right]^2 = f(\xi)$$

can be written

$$(4.2) \quad \left[ \pi i \phi(\zeta) + \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} \right] \cdot \left[ -\pi i \phi(\zeta) + \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} \right] = f(\zeta) .$$

We will solve this equation by using Carleman's method. Let us introduce the function

$$(4.3) \quad F(z) = \int_L \frac{\phi(\tau) d\tau}{\tau - z} ,$$

a function of the complex variable  $z = x + iy$ , which is analytic for  $z$  not on  $L$ , a simple smooth arc directed from  $z = -1$  to  $z = 1$ . If we let  $z$  approach a point  $\zeta$  on  $L$  from the positive side of  $L$ , and then the negative side of  $L$ ; the limit values of  $F(z)$  are

$$(4.4) \quad F^+(\zeta) = \pi i \phi(\zeta) + \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta}$$

$$(4.5) \quad F^-(\zeta) = -\pi i \phi(\zeta) + \int_L \frac{\phi(\tau) d\tau}{\tau - \zeta} .$$

Equation (4.2) says that these limit values must satisfy the barrier equation

$$(4.6) \quad F^+(\zeta) \cdot F^-(\zeta) = f(\zeta)$$

where  $\zeta$  is not an endpoint of  $L$ . The function  $F(z)$  which satisfies this equation and has the properties implied by the representation (4.3) provides the solution of the integral equation because the subtraction of the Plemelj formulas (4.4) and (4.5) gives

$$(4.7) \quad \phi(\zeta) = \frac{1}{2\pi i} [F^+(\zeta) - F^-(\zeta)] .$$

If we write

$$(4.8) \quad F(z) = \frac{e^{G(z)}}{z + i\sqrt{1-z^2}}$$

then (4.6) implies that  $G(z)$  must satisfy

$$(4.9) \quad G^+(\zeta) + G^-(\zeta) = \ln f(\zeta) + 2\pi i n$$

where  $n$  is an integer. As can be seen by using the Plemelj formulas, the function

$$(4.10) \quad G_1(z) = \frac{\sqrt{1-z^2}}{2\pi i} \int_L \frac{\ln f(\tau) d\tau}{(\sqrt{1-\tau^2})^+(\tau-z)} + \pi i n$$

is a particular solution of (4.9). By  $\sqrt{1-z^2}$  we mean here, and in what follows, the branch of the two valued function  $(1-z^2)^{1/2}$  which is analytic for any finite  $z$  not on  $L$ , and such that

$$\lim_{z \rightarrow \infty} \frac{\sqrt{1-z^2}}{z} = -1.$$

The function  $G_1(z)$  is analytic for  $z$  not on  $L$  and since

$$\lim_{z \rightarrow \infty} G_1(z) = \text{const.},$$

we have

$$\lim_{z \rightarrow \infty} \frac{e^{G_1(z)}}{z + i\sqrt{1-z^2}} = 0.$$

The general solution of (4.9) is

$$G(z) = G_1(z) + \sqrt{1-z^2} p(z)$$

where  $p(z)$  must satisfy



$$(4.11) \quad p^+(\zeta) - p^-(\zeta) = 0.$$

The function  $p(z)$  must be taken so that the properties of  $\phi(z)/z + i\sqrt{1-z^2}$  match those of

$$F(z) = \int_L \frac{\phi(\tau)d\tau}{\tau - z}.$$

This function is analytic for  $z$  not on  $L$  and it vanishes like  $c_0/z$  as  $z \rightarrow \infty$ , provided  $c_0 = \int_L \phi(\tau)d\tau \neq 0$ . Furthermore, in accordance with the assumptions about  $\phi(\tau)$  admitted in the introduction, the behavior of  $F(z)$  in the neighborhood of an endpoint  $\alpha$  of  $L$  is such that  $\lim_{z \rightarrow \alpha} (\alpha - z)F(z) = 0$ ; and the limit values  $F^+(\zeta)$ ,  $F^-(\zeta)$  must satisfy a uniform Hölder condition. These properties and the condition (4.11) imply that  $p(z)$  must be analytic everywhere and it must vanish at infinity, i.e.,  $p(z) = 0$ .

We have now found that

$$(4.12) \quad F(z) = \pm (z - i\sqrt{1-z^2}) \cdot \exp \left\{ \frac{\sqrt{1-z^2}}{2\pi i} \int_L \frac{\ln f(\tau)d\tau}{(\sqrt{1-\tau^2})^{\pm}(\tau-z)} \right\}.$$

This gives

$$(4.13) \quad F^+(\zeta) = \pm (\zeta - i\sqrt{1-\zeta^2}) \sqrt{f(\zeta)} \cdot \exp \left\{ \frac{\sqrt{1-\zeta^2}}{2\pi i} \int_L \frac{\ln f(\tau)d\tau}{\sqrt{1-\tau^2}(\tau-\zeta)} \right\}$$

$$(4.14) \quad F^-(\zeta) = \pm (\zeta + i\sqrt{1-\zeta^2}) \sqrt{f(\zeta)} \cdot \exp \left\{ -\frac{\sqrt{1-\zeta^2}}{2\pi i} \int_L \frac{\ln f(\tau)d\tau}{\sqrt{1-\tau^2}(\tau-\zeta)} \right\}$$

The solution of (4.1) is obtained by subtracting (4.14) from (4.13). It is

$$(4.15) \quad \phi(\zeta) = \frac{1}{2\pi i} [F^+(\zeta) - F^-(\zeta)]$$

$$= \pm \frac{\sqrt{1-\zeta^2}}{\pi} \left\{ \begin{aligned} &\zeta \sin \left[ \frac{\sqrt{1-\zeta^2}}{2\pi} \int_L \frac{\ln f(\tau) d\tau}{\sqrt{1-\tau^2}(\tau-\zeta)} \right] \\ &+ \sqrt{1-\zeta^2} \cos \left[ \frac{\sqrt{1-\zeta^2}}{2\pi} \int_L \frac{\ln f(\tau) d\tau}{\sqrt{1-\tau^2}(\tau-\zeta)} \right] \end{aligned} \right\}.$$

It will be noticed that in the above analysis we have assumed that  $F(z) = \int_L \frac{\phi(\tau) d\tau}{\tau - z}$  vanishes like  $c_0/z$ ,  $c_0 \neq 0$ , as  $z \rightarrow \infty$ . If this is not the case, that is, if for example

$$(4.16) \quad c_0 = - \int_L \phi(\tau) d\tau = 0$$

while

$$(4.17) \quad c_1 = - \int_L \tau \phi(\tau) d\tau = 0$$

then  $F(z)$  vanishes like  $c_1/z^2$ ,  $c_1 \neq 0$ , as  $z \rightarrow \infty$  and the solution of (4.6) as we have given it has to be adjusted. However, it is clear from the above analysis that if (4.16) and (4.17) prevail then the adjustment is easily made by taking  $F(z) = e^{G(z)}/(z + i\sqrt{1-z^2})^2$  instead of (4.8), which leads to

$$(4.18) \quad F(z) = \pm [z - i\sqrt{1-z^2}]^2 \exp \left\{ \frac{\sqrt{1-z^2}}{2\pi i} \int_L \frac{\ln f(\tau) d\tau}{(\sqrt{1-\tau^2})^+ (\tau-z)} \right\}.$$

### 5. Some Non-linear Problems in Potential Theory

The equations we have analyzed can be identified with certain problems in potential theory. For example, consider the problem of finding  $\psi(x,y)$  such that

$$\psi_{xx}(x,y) + \psi_{yy}(x,y) = 0, \quad y < 0,$$

$$\psi_y(x,0) = 0, \quad |x| > 1$$

and subject to an additional condition on  $y = 0$ ,  $|x| < 1$  which is specified below. Let us suppose that  $\psi_y(x,0)$ ,  $|x| < 1$ , satisfies the conditions imposed on  $\phi(t)$  in Section 1, and that each of  $\psi_x(x,y)$  and  $\psi_y(x,y)$  vanishes as  $z = x+iy \rightarrow \infty$ . The harmonic function  $\psi_y(x,y)$  can then be written

$$\begin{aligned} \psi_y(x,y) &= -\frac{1}{\pi} \int_{-1}^1 \frac{y\psi_y(t,0)dt}{(t-x)^2 + y^2} \\ &= -\frac{1}{2\pi} \frac{\partial}{\partial y} \int_{-1}^1 \ln[(t-x)^2 + y^2] \psi_y(t,0)dt \end{aligned}$$

from which

$$\psi_{xy}(x,y) = \frac{1}{\pi} \frac{\partial}{\partial y} \int_{-1}^1 \frac{(t-x)\psi_y(t,0)dt}{(t-x)^2 + y^2}$$

and by integration

$$\psi_x(x,y) = \frac{1}{\pi} \int_{-1}^1 \frac{(t-x)\psi_y(t,0)dt}{(t-x)^2 + y^2}$$

$$\psi_x(x,0) = \frac{1}{\pi} \int_{-1}^1 \frac{\psi_y(t,0)dt}{t-x}.$$

Now if we impose the additional condition

$$I \quad \frac{\lambda}{\pi} \psi_y(x,0) + \psi_x(x,0) \psi_y(x,0) = \frac{f(x)}{\pi}, \quad |x| < 1$$

then  $\phi(t) = \psi_y(t,0)$ ,  $|t| < 1$ , must satisfy

$$\lambda \phi(x) + \phi(x) \int_{-1}^1 \frac{\phi(t) dt}{t-x} = f(x), \quad |x| < 1.$$

If instead of I we impose

$$II \quad \psi_y^2(x,0) - \psi_x^2(x,0) = \frac{f(x)}{\pi^2}, \quad |x| < 1$$

then  $f(t) = \psi_y(t,0)$ , must satisfy

$$\pi^2 \phi^2(x) - \left[ \int_{-1}^1 \frac{\phi(t) dt}{t-x} \right]^2 = f(x), \quad |x| < 1.$$

Finally, if the additional condition is

$$III \quad \psi_y^2(x,0) + \psi_x^2(x,0) = \frac{f(x)}{\pi^2}, \quad |x| < 1$$

then  $\phi(t) = \psi_y(t,0)$ , must satisfy

$$\pi^2 \phi^2(x) + \left[ \int_{-1}^1 \frac{\phi(t) dt}{t-x} \right]^2 = f(x), \quad |x| < 1.$$

We proceed to show that problem III can be solved for domains more general than the half plane. For the half plane problem we can write

$$\psi(x,y) = \Re F(z)$$

and then

$$\mathcal{F}'(z) = \mathcal{R}\mathcal{F}'(z) + i\mathcal{I}\mathcal{F}'(z) = \psi_x(x,y) - i\psi_y(x,y)$$

$$= \frac{1}{\pi} \mathcal{R} \int_{-1}^1 \frac{\phi(t)dt}{t-z} + \frac{1}{\pi} \mathcal{I} \int_{-1}^1 \frac{\phi(t)dt}{t-z}$$

which is the same as

$$(5.1) \quad \mathcal{F}'(z) = \frac{1}{\pi} \int_{-1}^1 \frac{\phi(t)dt}{t-z}.$$

The integral on the right hand side of (5.1) has been determined in connection with the problem of solving the integral equation III, in Section 4, and we have

$$(5.2) \quad \mathcal{F}'(z) = \pm \frac{(z-1\sqrt{1-z^2})^\gamma}{\pi} \exp \left\{ \frac{\sqrt{1-z^2}}{2\pi i} \int_{-1}^1 \frac{\ln f(t)dt}{(\sqrt{1-t^2})^+(t-z)} \right\}$$

where  $\gamma$  is 1, or 2. The function  $\psi(x,y)$  can be obtained from (5.2) by an integration.

Let  $z = m(\zeta) = m(\xi + i\eta)$  be a function which maps the domain  $D$ , with boundary  $C$  in the  $\zeta$ -plane, conformally into the lower half of the  $z$ -plane. Let the image of  $C_1$  a part of  $C$ , be the segment  $y = 0$ ,  $|x| < 1$ ; and let the image of  $C_2$ , the remaining part of  $C$ , be  $y = 0$ ,  $|x| > 1$ . Under this mapping,  $\mu(x,y)$  is transformed into

$$\psi(x,y) = \underline{\Psi}(\xi,\eta) = \mathcal{R}\mathcal{F}[m(\zeta)],$$

a function harmonic in  $D$ .

Let  $s$  be the arc length measured along  $C$  from say the initial point of  $C_1$ . If  $\zeta = \xi(s) + i\eta(s)$  is a point on  $C$ , then the normal derivative of  $\underline{\Psi}(\xi,\eta)$  at the boundary is

$$\begin{aligned}
 (5.3) \quad \frac{\partial \underline{\Psi}(\xi, \eta)}{\partial n} &= \frac{\partial}{\partial n} \mathcal{R} \mathcal{F}[m(\sigma)] \\
 &= \frac{\partial}{\partial s} \mathcal{I} \mathcal{F}[m(\sigma)] \\
 &= \mathcal{I} \frac{\partial m(\sigma)}{\partial s} \mathcal{F}'[m(\sigma)] .
 \end{aligned}$$

If  $\sigma$  is on  $C_2$ , then  $\frac{dm(\sigma)}{ds} = \frac{dx}{ds}$  is real and  $\mathcal{F}'[m(\sigma)]$  is real.

Hence for  $\sigma$  on  $C_2$

$$\underline{\Psi}_n(\xi, \eta) = 0 .$$

The tangential derivative of  $\underline{\Psi}(\xi, \eta)$  at the boundary is

$$(5.4) \quad \frac{\partial \underline{\Psi}(\xi, \eta)}{\partial s} = \mathcal{R} \frac{\partial m(\sigma)}{\partial s} \mathcal{F}'[m(\sigma)] .$$

From (5.3) and (5.4) we see that for  $\sigma$  on  $C_1$

$$\begin{aligned}
 \underline{\Psi}_n^2(\xi, \eta) + \underline{\Psi}_s^2(\xi, \eta) &= \left| \frac{dm(\sigma)}{ds} \right|^2 \left| \mathcal{F}'[m(\sigma)] \right|^2 \\
 &= \left| \frac{dm(\sigma)}{ds} \right|^2 \left[ \{ \mathcal{R} \mathcal{F}'[m(\sigma)] \}^2 + \{ \mathcal{I} \mathcal{F}'[m(\sigma)] \}^2 \right] \\
 &= \left| \frac{dm(\sigma)}{ds} \right|^2 \cdot \frac{f[m(\sigma)]}{\pi^2} .
 \end{aligned}$$

Therefore if we take

$$h(s) = \left| \frac{dm(\sigma)}{ds} \right|^2 \frac{f[m(\sigma)]}{\pi^2}$$

as prescribed, then

$$\underline{\Psi}(\xi, \eta) = \mathcal{R} \mathcal{F}[m(\zeta)]$$

is harmonic in  $D$ , with normal derivative equal to zero along  $C_2$ , and with

$$\psi_n^2(\xi, \eta) + \psi_s^2(\xi, \eta) = h(s)$$

prescribed along  $C_1$ . Thus the solution of problem III for the domain D can be found by substituting  $z = m(\zeta)$ ,  $t = m(\sigma)$  and

$$f(t) = f[m(\sigma)] = \frac{\pi^2 h(s)}{\left| \frac{dm(\sigma)}{ds} \right|^2}$$

in (5.2). The substitution gives

$$(5.5) \quad \frac{d\mathcal{F}[m(\zeta)]}{d\zeta} = \pm \frac{m'(\zeta)}{\pi} \left[ m(\zeta) - i\sqrt{1-m^2(\zeta)} \right]^\gamma \cdot \exp \left\{ -\frac{\sqrt{1-m^2(\zeta)}}{2\pi i} \int_{C_1} \frac{\ln \left\{ \left| \frac{dm(\sigma)}{ds} \right|^2 \cdot h(s) \right\} \cdot \frac{dm(\sigma)}{ds} ds}{\sqrt{1-m^2(\sigma)} [m(\sigma) - m(\zeta)]} \right\}$$

and then  $\psi(\xi, \eta)$  can be obtained by integration.

The solution of problem III for particular domains and special boundary values of the gradient is not new. For example, it is well known that the method of conformal mapping can be used to find a potential function  $\psi(\xi, \eta)$  such that

$$\psi_{\xi\xi}(\xi, \eta) + \psi_{\eta\eta}(\xi, \eta) = 0, \quad -\infty < \xi < \infty; \quad -1 < \eta < 1;$$

$$\psi_n(\xi, \eta) = 0 \quad \eta = -1, 1; \quad -\infty < \xi < 0;$$

$$\psi_\xi^2(\xi, \eta) + \psi_\eta^2(\xi, \eta) = \text{const.} \quad \eta = -1, 1; \quad 0 < \xi < \infty.$$

This problem, or its equivalent, occurs in the hydrodynamical theory of jets. However, the author has not seen the formula (5.5) in the literature and it is probable that there are several problems in field or flow theory for which it would be advantageous to use (5.5) directly.

In order to see a simple application of (5.5) consider the problem of finding  $\Psi(\xi, \eta)$  such that

$$\begin{aligned}
 (5.6) \quad & \Psi_{\xi\xi}(\xi, \eta) + \Psi_{\eta\eta}(\xi, \eta) = 0, & -\infty < \xi < \infty; & 0 < \eta < 1 \\
 & \Psi_{\eta}(\xi, 0) = 0, & -\infty < \xi < \infty \\
 & \Psi_{\xi}^2(\xi, 1) + \Psi_{\eta}^2(\xi, 1) = h(\xi), & -\infty < \xi < \infty
 \end{aligned}$$

The function

$$(5.7) \quad z = x + iy = \coth \frac{\pi}{2} (\xi + i\eta) = \coth \frac{\pi}{2} \zeta = m(\zeta)$$

maps the infinite strip into the lower half of the  $z$ -plane and the segment  $\mathcal{I} z = 0; |R(z)| < 1$  is the image of  $C_1: \zeta = \xi + 1; -\infty < \xi < \infty$ . From (5.7) we find

$$m'(\zeta) = -\frac{\pi}{2} \operatorname{csch}^2 \frac{\pi}{2} \zeta$$

$$\sqrt{1 - m^2(\zeta)} = \sqrt{1 - \coth^2 \frac{\pi}{2} \zeta} = -1 \operatorname{csch} \frac{\pi}{2} \zeta$$

$$m(\sigma) = m(\lambda + i) = \tanh \frac{\pi}{2} \lambda$$

$$\sqrt{1 - m^2(\sigma)} = \sqrt{1 - \tanh^2 \frac{\pi}{2} \lambda} = \operatorname{sech} \frac{\pi}{2} \lambda$$

$$\frac{dm(\sigma)}{ds} = \frac{d}{d\lambda} \left( \tanh \frac{\pi}{2} \lambda \right) \frac{d\lambda}{ds}, \quad \frac{d\lambda}{ds} = -1$$

$$= -\frac{\pi}{2} \operatorname{sech}^2 \frac{\pi}{2} \lambda,$$

$$\frac{1}{m(\sigma) - m(\zeta)} = -\frac{\cosh \frac{\pi}{2} \lambda \cdot \sinh \frac{\pi}{2} \zeta}{\cosh \frac{\pi}{2} (\lambda - \zeta)}.$$



The substitution of these quantities in (5.5) gives

$$(5.8) \quad \frac{d}{d\zeta} \mathcal{F}[m(s)] = \pm \frac{1}{2 \sinh^2 \frac{\pi}{2} \zeta} \left[ \frac{\cosh \frac{\pi}{2} \zeta - 1}{\sinh \frac{\pi}{2} \zeta} \right]^\gamma \\ \cdot \exp \left\{ \frac{1}{4} \int_{-\infty}^{\infty} \frac{\ln[4 \cdot \cosh^4 \frac{\pi}{2} \lambda \cdot h(\lambda)] d\lambda}{\cosh \frac{\pi}{2} (\lambda - \zeta)} \right\}$$

which if we use

$$(5.9) \quad \frac{1}{4} \int_{-\infty}^{\infty} \frac{\ln[4 \cosh^4 \frac{\pi}{2} \lambda] d\lambda}{\cosh \frac{\pi}{2} (\lambda - \zeta)} \\ = \ln 2 + 2 \ln[1 + \cosh \frac{\pi}{2} \zeta] + 2\pi i \eta$$

becomes

$$(5.10) \quad \frac{d}{d\zeta} \mathcal{F}[m(\zeta)] = \pm \frac{2}{2 \sinh^2 \frac{\pi}{2} \zeta} \left[ \frac{\cosh \frac{\pi}{2} \zeta - 1}{\sinh \frac{\pi}{2} \zeta} \right]^\gamma \cdot \left[ \cosh \frac{\pi}{2} \zeta + 1 \right]^2 \\ \cdot \exp \frac{1}{4} \int_{-\infty}^{\infty} \frac{h(\lambda) d\lambda}{\cosh \frac{\pi}{2} (\lambda - \zeta)}.$$

If we choose  $\gamma = 2$  so as to avoid a singularity at  $\zeta = 0$  we finally have

$$(5.11) \quad \frac{d}{d\zeta} \mathcal{F}[m(\zeta)] = \pm \exp \left\{ \frac{1}{4} \int_{-\infty}^{\infty} \frac{h(\lambda) d\lambda}{\cosh \frac{\pi}{2} (\lambda - \zeta)} \right\}$$

which determines the derivatives of  $\Psi(\xi, \eta)$ . That is, if

$$\Psi(\xi, \eta) = \mathcal{R} M(\zeta)$$

is required to satisfy (5.6) then

$$M'(\zeta) = \pm \exp \left\{ \frac{1}{4} \int_{-\infty}^{\infty} \frac{h(\lambda) d\lambda}{\cosh \frac{\pi}{2} (\lambda - \zeta)} \right\}.$$

## 6. Some Non-linear Systems

In a recent paper, A. Beurling [5] proved that if  $D$  is a region in Euclidean  $n$ -space then the non-linear system

$$(6.1) \quad \begin{aligned} \phi(x) \int_D k(t,x) \psi(t) dt &= f(x) \\ \psi(x) \int_D k(x,t) \phi(t) dt &= g(x) \end{aligned}$$

possesses a unique solution when the kernel  $k(t,x)$  and the prescribed functions  $f(x)$ ,  $g(x)$  are suitably restricted. Since Beurling's analysis does not cover the following singular one-dimensional case

$$(6.2) \quad \begin{aligned} \phi(x) \int_{-1}^1 \frac{\psi(t) dt}{t-x} &= f(x) \\ \psi(x) \int_{-1}^1 \frac{\phi(t) dt}{t-x} &= g(x) \end{aligned} \quad -1 < x < 1$$

and since this system is related to certain problems in potential theory; it is of some interest to see that (6.2) can be reduced to a linear system by using the method of Section 2.

We suppose that each of the unknown functions belongs to the class  $\mathcal{N}_1$ , defined in the introduction, and we suppose that each of the prescribed functions  $f(x)$ ,  $g(x)$  belongs to the class  $\mathcal{N}_2$ . Then, by using the Hardy-Poincaré-Bertrand formula as shown in Section 2 it is easy to verify that

$$\begin{aligned}
 (6.3) \quad & \frac{1}{1-\xi^2} \int_{-1}^1 \frac{(1-x^2)}{x-\xi} \left[ \phi(x) \int_{-1}^1 \frac{\psi(t)dt}{t-x} + \psi(x) \int_{-1}^1 \frac{\phi(t)dt}{t-x} \right] dx - \frac{a_0 b_0}{1-\xi^2} \\
 & = -\pi^2 \phi(\xi) \psi(\xi) + \int_{-1}^1 \frac{\psi(t)dt}{t-\xi} \cdot \int_{-1}^1 \frac{\phi(t)dt}{t-\xi}
 \end{aligned}$$

where

$$a_0 = \int_{-1}^1 \phi(t)dt$$

$$b_0 = \int_{-1}^1 \psi(t)dt.$$

If (6.2) possesses a solution, then (6.2) and (6.3) imply

$$\begin{aligned}
 (6.4) \quad & \frac{1}{1-\xi^2} \int_{-1}^1 \frac{(1-x^2)[f(x)+g(x)]dx}{x-\xi} - \frac{a_0 b_0}{1-\xi^2} \\
 & = -\pi^2 \phi(\xi) \psi(\xi) + \int_{-1}^1 \frac{\psi(t)dt}{t-\xi} \cdot \int_{-1}^1 \frac{\phi(t)dt}{t-\xi}
 \end{aligned}$$

or after multiplying by  $\phi(\xi)\psi(\xi)$

$$\begin{aligned}
 (6.5) \quad & \pi^2 \phi^2(\xi) \psi^2(\xi) + \frac{\phi(\xi)\psi(\xi)}{1-\xi^2} \left[ \int_{-1}^1 \frac{(1-x^2)[f(x)+g(x)]dx}{x-\xi} - a_0 b_0 \right] \\
 & - \phi(\xi) \int_{-1}^1 \frac{\psi(t)dt}{t-\xi} \cdot \psi(\xi) \int_{-1}^1 \frac{\phi(t)dt}{t-\xi} = 0.
 \end{aligned}$$

Using (6.2) again, equation (6.5) becomes

$$(6.6) \quad \pi^2 \phi^2(\xi) \psi^2(\xi) + \frac{\phi(\xi)\psi(\xi)}{1-\xi^2} F(\xi) - f(\xi)g(\xi) = 0$$

where

$$F(\xi) = \int_{-1}^1 \frac{(1-x^2)[f(x)+g(x)]dx}{x-\xi} - a_0 b_0.$$

Equation (6.6) fixes the product  $\phi(\xi)\psi(\xi)$ . We find

$$(6.7) \quad \phi(\xi)\psi(\xi) = k(\xi) = - \frac{F(\xi) \pm \sqrt{F^2(\xi) + 4\pi^2 f(\xi)g(\xi)(1-\xi^2)^2}}{2\pi^2(1-\xi^2)}.$$

With this product known the system (6.2) can be written

$$(6.8) \quad \int_{-1}^1 \frac{\psi(t)dt}{t-x} = \frac{f(x)}{k(x)} \cdot \psi(x),$$

$$\int_{-1}^1 \frac{\phi(t)dt}{t-x} = \frac{g(x)}{k(x)} \cdot \phi(x).$$

The last equations can now be solved by Carleman's method [4].

Notice that if each of  $f(x)$  and  $g(x)$  is integrable and we require each of the unknown functions to satisfy the condition (B) of Section 2 then

$$\begin{aligned} \int_{-1}^1 f(x)dx &= \int_{-1}^1 \phi(x) \int_{-1}^1 \frac{\psi(t)dt dx}{t-x} = - \int_{-1}^1 \psi(t) \int_{-1}^1 \frac{\phi(x)dx dt}{x-t} \\ &= - \int_{-1}^1 g(t)dt \end{aligned}$$

or

$$(6.9) \quad \int_{-1}^1 [f(x) + g(x)]dx = 0.$$

Also,

$$\begin{aligned} \int_{-1}^1 xf(x)dx &= - \int_{-1}^1 \psi(t) \int_{-1}^1 \frac{(x-t+t)}{x-t} \phi(x)dx dt \\ &= - \int_{-1}^1 \psi(t)dt \cdot \int_{-1}^1 \phi(x)dx - \int_{-1}^1 t\psi(t) \int_{-1}^1 \frac{\phi(x)dx dt}{x-t} \end{aligned}$$

which determines  $a_0 b_0$ , that is,

$$-a_0 b_0 = \int_{-1}^1 x[f(x) + g(x)]dx .$$

Hence for solutions which satisfy (B),  $F(\xi)/(1-\xi^2)$  becomes

$$\frac{F(\xi)}{1-\xi^2} = \int_{-1}^1 \frac{[f(x) + g(x)]dx}{x-\xi} - \frac{\xi}{1-\xi^2} \int_{-1}^1 [f(x) + g(x)]dx$$

and since the second term on the right is zero by (6.9) we have

$$\frac{F(\xi)}{1-\xi^2} = \int_{-1}^1 \frac{[f(x) + g(x)]dx}{x-\xi} .$$

We remark in passing that the system

$$\begin{aligned} \phi_1(x) \int_{-1}^1 \frac{\psi_1(t)dt}{t+x} &= f_1(x) \\ \psi_1(x) \int_{-1}^1 \frac{\phi_1(t)dt}{t+x} &= g_1(x) \end{aligned} \quad -1 < x < 1$$

can be reduced to (6.2) by using the substitutions

$$\begin{aligned} \phi_1(t) &= \phi(-t) , & f_1(x) &= f(-x) , \\ \psi_1(t) &= \psi(t) , & g_1(x) &= -g(x) . \end{aligned}$$

The systems noted above are not, of course, the only non-linear systems that can be solved by using the technique of Section 2. The Hardy-Poincaré-Bertrand formula can be used to linearize a number of other systems in which the non-linearity is due to the presence of a product in which an unknown function is

multiplied by an unknown Hilbert transform or an unknown Stieltjes transform. As two additional examples we cite the systems

$$\phi(x) \int_{-1}^1 \frac{\psi(t)dt}{t-x} = f(x) + \lambda_1 \phi(x)$$

$$\psi(x) \int_{-1}^1 \frac{\phi(t)dt}{t-x} = g(x) + \lambda_2 \psi(x)$$

and

$$-\phi(x) \int_0^1 \frac{\phi(t)dt}{t-x} + \psi(x) \int_0^1 \frac{\psi(t)dt}{t+x} = f_1(x)$$

$$\phi(x) \int_0^1 \frac{\psi(t)dt}{t-x} - \psi(x) \int_0^1 \frac{\phi(t)dt}{t-x} = f_2(x) .$$

The last system plays an important role in the theory of radiative transfer.

### References

- [1] Radiative Transfer by S. Chandrasekhar. Oxford University Press, (1950).
- [2] Basic Methods in Transfer Problems by V. Kourganoff; I. W. Busbridge (Coll.). Oxford University Press, (1952).
- [3] Fox, C., A Solution of Chandrasekhar's Integral Equation, Trans. Amer. Math. Soc. 99, (1961), 285-291.
- [4] Singular Integral Equations by N. I. Muskhelishvili. P. Noordhoff N.V., Groningen, Holland (1953).
- [5] Beurling, A., An Automorphism of Product Measures, Ann. of Math. (2) 72 (1960), 189-200.

## Approved Distribution List

N-1

Chief of Naval Research Department of the Navy Washington 25, D.C. Attn: Code 438	(3)	Chief, Bureau of Aeronautics Department of the Navy Washington 25, D.C. Attn: Research Division	(1)
Commanding Officer Office of Naval Research Branch Office The John Crerar Library Bldg. 86 East Randolph Street Chicago 1, Illinois	(1)	Chief, Bureau of Ordnance Department of the Navy Washington 25, D.C. Attn: Research and Development Division	(1)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, N.Y.	(2)	Office of Ordnance Research Department of the Army Washington 25, D.C.	(1)
Commanding Officer Office of Naval Research Branch Office 1000 Geary Street San Francisco 24, Calif.	(1)	Commander Air Research and Development Command Office of Scientific Research P.O. Box 1395 Baltimore 18, Md. Attn: Fluid Mechanics Div.	(1)
Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena 1, Calif.	(1)	Director of Research National Advisory Committee for Aeronautics 1724 F Street, Northwest Washington 25, D.C.	(1)
Commanding Officer Office of Naval Research Navy No. 100, Fleet Post Office New York, New York	(5)	Director Langley Aeronautical Laboratory National Advisory Committee for Aeronautics Langley Field, Virginia	(1)
Director Naval Research Laboratory Washington 25, D.C. Attn: Code 2021	(6)	Director National Bureau of Standards Washington 25, D.C. Attn: Fluid Mechanics Section	(1)
Documents Service Center Armed Services Technical Information Agency Knott Building Dayton 2, Ohio	(5)	Professor R. Courant Institute of Mathematical Sciences, N.Y.U. 25 Waverly Place New York 3, New York	(1)
Professor W.R. Sears Director Graduate School of Aeronautical Engineering Cornell University Ithaca, New York	(1)	Professor G. Kuerti Department of Mechanical Engineering Case Institute of Technology Cleveland, Ohio	(1)



## Distribution List (Cont.)

N-11

Chief, Bureau of Ships  
Department of the Navy  
Washington 25, D.C.  
Attn: Research Division (1)  
Code 420 Preliminary Design (1)

Commander  
Naval Ordnance Test Station  
3202 E. Foothill Blvd.  
Pasadena, Calif. (1)

Commanding Officer and Director  
David Taylor Model Basin  
Washington 7, D.C.  
Attn: Hydromechanics Lab (1)  
Hydrodynamics Div. (1)  
Library (1)

California Institute of  
Technology  
Hydrodynamic Laboratory  
Pasadena 4, Calif. (1)

Professor A.T. Ippen  
Hydrodynamics Laboratory  
Massachusetts Institute  
of Technology  
Cambridge 39, Mass. (1)

Dr. Hunter Rouse, Director  
Iowa Institute of Hydraulic  
Research  
State University of Iowa  
Iowa City, Iowa (1)

Stevens Institute of Technology  
Experimental Towing Tank  
711 Hudson St.  
Hoboken, New Jersey (1)

Dr. L.G. Straub  
St. Anthony Falls Hydraulic  
Laboratory  
University of Minnesota  
Minneapolis 14, Minn. (1)

Dr. G.H. Hickox  
Engineering Experiment Station  
University of Tennessee  
Knoxville, Tenn. (1)

Chief, of Naval Research  
Department of the Navy  
Washington 25, D.C.  
Attn: Code 416 (1)  
Code 460 (1)

Chief, Bureau of Yards  
and Docks  
Department of the Navy  
Washington 25, D.C.  
Attn: Research Division (1)

Commanding Officer and Director  
David Taylor Model Basin  
Washington 7, D.C.  
Attn: Ship Division (1)

Hydrographer  
Department of the Navy  
Washington 25, D.C. (1)

Director  
Waterways Experiment Station  
Box 631  
Vicksburg, Mississippi (1)

Office of the Chief of Engineer  
Department of the Army  
Gravelly Point  
Washington 25, D.C. (1)

Beach Erosion Board  
U.S. Army Corps of Engineers  
Washington 25, D.C. (1)

Commissioner  
Bureau of Reclamation  
Washington 25, D.C. (1)

Dr. G.H. Keulegan  
National Hydraulic Laboratory  
National Bureau of Standards  
Washington 25, D.C. (1)

Brown University  
Graduate Division of Applied  
Mathematics  
Providence 12, Rhode Island (1)

California Institute of  
Technology  
Hydrodynamics Laboratory  
Pasadena 4, Calif.  
Attn: Professor M.S. Plesset (1)  
Professor V.A. Vanoni (1)

## Distribution List (Cont.)

N-111

Mr. C.A. Gongwer Aerojet General Corporation 6352 N. Irwindale Avenue Azusa, Calif.	(1)	Director, USAF Project RAND Via: Air Force Liaison Office The RAND Corporation 1700 Main Street Santa Monica, Calif.	(1)
Professor M.L. Albertson Department of Civil Engineering Colorado A. + M. College Fort Collins, Colorado	(1)	Attn: Library Commanding Officer NROTC and Naval Administrative Unit Massachusetts Institute of Tech. Cambridge 39, Massachusetts	(1)
Professor G. Birkhoff Department of Mathematics Harvard University Cambridge 38, Mass.	(1)	Commanding Officer and Director U.S. Navy Mine and Defense Laboratory Panama City, Florida	(1)
Massachusetts Institute of Technology Department of Naval Architecture Cambridge 39, Mass.	(1)		
Dr. R.R. Revelle Scripps Institute of Oceanography La Jolla, California	(1)		
Stanford University Applied Mathematics and Statistics Laboratory Stanford, California	(1)		
Professor J.W. Johnson Fluid Mechanics Laboratory University of California Berkeley 4, California	(1)		
Professor H.A. Einstein Department of Engineering University of California Berkeley 4, Calif.	(1)		
Dean K.E. Schoenherr College of Engineering University of Notre Dame Notre Dame, Indiana	(1)		
Director Woods Hole Oceanographic Institute Woods Hole, Mass.	(1)		
Hydraulics Laboratory Michigan State College East Lansing, Michigan Attn: Professor H.R. Henry	(1)		